```
? show files
File 155:MEDLINE(R) 1966-2004/Jun W2
          (c) format only 2004 The Dialog Corp.
       5:Biosis Previews(R) 1969-2004/Jun W4
          (c) 2004 BIOSIS
      35:Dissertation Abs Online 1861-2004/May
File
          (c) 2004 ProQuest Info&Learning
      47: Gale Group Magazine DB(TM) 1959-2004/Jun 28
File
          (c) 2004 The Gale group
      71:ELSEVIER BIOBASE 1994-2004/Jun W3
File
          (c) 2004 Elsevier Science B.V.
      73:EMBASE 1974-2004/Jun W3
File
          (c) 2004 Elsevier Science B.V.
File 144: Pascal 1973-2004/Jun W3
          (c) 2004 INIST/CNRS
File 351:Derwent WPI 1963-2004/UD,UM &UP=200440
          (c) 2004 Thomson Derwent
File 357: Derwent Biotech Res. 1982-2004/Jun W4
          (c) 2004 Thomson Derwent & ISI
File 358: Current BioTech Abs 1983-2004/May
           (c) 2004 DECHEMA
? ds s2-
                 Description
Set
        Items
                 (PROTEIN OR PEPTIDE? ? OR POLYPEPTIDE? ?) (5N) (DESIGN OR PR-
S2
              EPARATION OR GENERATE OR GENERATION OR ENGINEER? OR MANUFACTU-
              R? OR PRODUC? OR SYNTH?)
                 (CAMD OR CAD OR COMPUTER(W) (AIDED OR ASSISTED) (2N) DESIGN)
S3
                 (BIND OR BINDING OR ATTACH?) (5N) TERMI?
S4
        73732
S5
           15
                 S2 AND S3 AND S4
            9
                 RD (unique items)
S6
            5
                 S6 NOT CALDESMON
S7
?
? t s7/3 ab/1-5
            (Item 1 from file: 155)
 7/AB/1
DIALOG(R) File 155: MEDLINE(R)
(c) format only 2004 The Dialog Corp. All rts. reserv.
10261806
            PMID: 7525561
  Identification of the regulatory domain of the mammalian multifunctional
protein CAD by the construction of an Escherichia coli hamster hybrid
carbamyl-phosphate synthetase.
  Liu X; Guy H I; Evans D R
  Department of Biochemistry, Wayne State University School of Medicine,
Detroit, Michigan 48201.
  Journal of biological chemistry (UNITED STATES)
                                                         Nov 4 1994, 269 (44)
 p27747-55, ISSN 0021-9258 Journal Code: 2985121R
  Contract/Grant No.: GM 47399; GM; NIGMS
  Document type: Journal Article
  Languages: ENGLISH
  Main Citation Owner: NLM
  Record type: Completed
Carbamyl-phosphate synthetases from different organisms have similar catalytic mechanisms and amino acid sequences, but their structural organization, sub-unit structure, and mode of regulation can be very
different. Escherichia coli carbamyl-phosphate synthetase (CPSase), a
monofunctional protein consisting of amido-transferase and
synthetase subunits, is allosterically inhibited by UMP and activated
by NH3, IMP, and ornithine. In contrast, mammalian CPSase II, part of the
```

```
168349 SEA FILE=HCAPLUS ABB=ON PLU=ON (?TERMIN? OR ?DOMAIN?)(L)(?BIN
L10
                D? OR ?ATTACH?)
         117189 SEA FILE=HCAPLUS ABB=ON PLU=ON L5(L)L10
L11
                                                  L9 AND L11
            771 SEA FILE=HCAPLUS ABB=ON PLU=ON
L12
                                                  L3 OR 'IN SILICO' OR GLASS
                                         PLU=ON
        1068575 SEA FILE=HCAPLUS ABB=ON
L13
                                                  L12 AND L13
L14 NOT L2
             10 SEA FILE=HCAPLUS ABB=ON
                                         PLU=ON
L14
             10 SEA FILE=HCAPLUS ABB=ON
                                          PLU=ON
L15
            287 SEA FILE=HCAPLUS ABB=ON PLU=ON ENGINEERED (L) PROTEIN(L) BINDIN
L23
                G(L) DOMAINS
                                          PLU=ON L23 AND L7
              8 SEA FILE=HCAPLUS ABB=ON
L24
                                          PLU=ON L24 NOT (L2 OR L15)
              7 SEA FILE=HCAPLUS ABB=ON
L25
                                          PLU=ON COMPUTER(W) (ASSISTED OR
           2882 SEA FILE=HCAPLUS ABB=ON
L26
                AIDED) (2W) DESIGN
            251 SEA FILE=HCAPLUS ABB=ON PLU=ON L26 AND L5
L27
                                          PLU=ON L27 AND L11
             16 SEA FILE=HCAPLUS ABB=ON
L28
                                          PLU=ON L28 NOT (L2 OR L15 OR L25)
             14 SEA FILE=HCAPLUS ABB=ON
L29
            277 SEA FILE=HCAPLUS ABB=ON PLU=ON COMPUTER(W) (ASSISTED OR
L30
                AIDED) (W) DRUG (W) DESIGN
              8 SEA FILE=HCAPLUS ABB=ON PLU=ON L29 NOT L30
L31
=>
=>
=> d ibib abs hitrn 131 1-8
L31 ANSWER 1 OF 8 HCAPLUS COPYRIGHT 2004 ACS on STN
                         2001:303055 HCAPLUS 135:101926
ACCESSION NUMBER:
DOCUMENT NUMBER:
                          Design of inhibitors of Ras-Raf interaction using a
TITLE:
                          computational combinatorial algorithm
                          Zeng, Jun; Nheu, Thao; Zorzet, Anna; Catimel, Bruno;
AUTHOR(S):
                          Nice, Ed; Maruta, Hiroshi; Burgess, Antony W.;
                          Treutlein, Herbert R.
                          Ludwig Institute for Cancer Research, Royal Melbourne
CORPORATE SOURCE:
                          Hospital, Parkville, 3050, Australia
                          Protein Engineering (2001), 14(1), 39-45
SOURCE:
                          CODEN: PRENE9; ISSN: 0269-2139
                          Oxford University Press
PUBLISHER:
DOCUMENT TYPE:
                          Journal
                          English
LANGUAGE:
     Drugs that inhibit important protein-protein
     interactions are hard to find either by screening or rational design, at
     least so far. Most drugs on the market that target proteins
     today are therefore aimed at well-defined binding pockets in
     proteins. While computer-aided design
     is widely used to facilitate the drug discovery process for binding
     pockets, its application to the design of inhibitors that target the
     protein surface initially seems to be limited because of the
     increased complexity of the task. Previously, we had started to develop a
     computational combinatorial design approach based on the well-known
     "multiple copy simultaneous search" (MCSS) procedure to tackle this
     problem. In order to identify sequence patterns of potential inhibitor peptides, a three-step procedure is employed: first, using MCSS,
     the locations of specific functional groups on the protein
     surface are identified; second, after constructing the peptide
     main chain based on the location of favorite locations of
     N-methylacetamide groups, functional groups corresponding to amino acid
     side chains are selected and connected to the main chain C\alpha atoms;
      finally, the peptides generated in the second step are aligned
```

and probabilities of amino acids at each position are calculated from the alignment scheme. Sequence patterns of potential inhibitors are determined

based on the propensities of amino acids at each $C\alpha$ position. Here